

10/511627

WEICKMANN & WEICKMANN

DT05 Rec'd PCT/PTO 18 OCT 2004

Patentanwälte  
European Patent Attorneys · European Trademark Attorneys

WEICKMANN & WEICKMANN, Postfach 860 820, 81635 München

European Patent Office

80298 Munich

DIPL.-ING. H. WEICKMANN (bis 31.1.01)  
DIPL.-ING. F. A. WEICKMANN  
DIPL.-CHEM. B. HUBER  
DR.-ING. H. LISKÄ  
DIPL.-PHYS. DR. J. PRECHTEL  
DIPL.-CHEM. DR. B. BOHM  
DIPL.-CHEM. DR. W. WEISS  
DIPL.-PHYS. DR. J. TIESMEYER  
DIPL.-PHYS. DR. M. HERZOG  
DIPL.-PHYS. B. RUTTENSBERGER  
DIPL.-PHYS. DR.-ING. V. JORDAN  
DIPL.-CHEM. DR. M. DEY  
DIPL.-FORSTW. DR. J. LACHNIT

IHR ZEICHEN / YOUR REF.

UNSER ZEICHEN / OUR REF.

DATUM / DATE

28050P WO/WWCGLd

2. Juli 2003

*Auth*

Application No./Patent No.:

PCT/EP 03/04650

Applicant/Proprietor:

DeveloGen Aktiengesellschaft für entwicklungsbiologische Forschung

Encl.

Sequence Listing (diskette) ✓

Sequence Listing (paper  
copy) (all 1-fold) ✓

New description pages 60 ✓  
and 61 ✓

In response to the official communication  
dated June 26, 2003.

Applicants herewith submit a sequence listing  
in computer-readable form as well as in the  
form of a paper copy.


It is stated that the information recorded on  
the data carrier is identical to the written  
sequence listing. It is further stated that the  
sequence listing does not include subject  
matter which goes beyond the content of the  
application as originally filed.

Postfach 860 820  
81635 München  
Kopernikusstraße 9  
81679 München  
Deutschland

Telefon (089) 45563 0  
(0700) WEICKMANN  
Telefax (089) 45563 999  
E-Mail email@weickmann.de  
Internet www.weickmann.de  
Vat-ID. Nr. DE 130 753 315

Hypovereinsbank München  
Konto 208 401 (BLZ 700 202 70)  
S.W.I.F.T.-Adresse HYVE DE MM  
Postbank München  
Konto 77 46-804 (BLZ 700 100 80)  
S.W.I.F.T.-Adresse PBNKDEFF 700

Furthermore, new description pages 60 and 61 are filed including the correct sequence numberings.



Dr. W. Weiss

temperature (preferably 22°C), 40 per cent humidity and a light / dark cycle of preferably 14 / 10 hours. The mice were fed a standard chow (for example, from ssniff Spezialitäten GmbH, order number ssniff M-Z V1126-000). For the fasting experiment ("fasted wild type mice"), wild type mice were starved for 48 h without food, but only water supplied ad libitum (see, for example, Schnetzler et al., (1993) J Clin Invest 92(1):272-280, Mizuno et al., (1996) Proc Natl Acad Sci U S A 93(8):3434-3438). Animals were sacrificed at an age of 6 to 8 weeks. The animal tissues were isolated according to standard procedures known to those skilled in the art, snap frozen in liquid nitrogen and stored at -80°C until needed.

RNA was isolated from mouse tissues using Trizol Reagent (for example, from Invitrogen, Karlsruhe, Germany) and further purified with the RNeasy Kit (for example, from Qiagen, Germany) in combination with an DNase-treatment according to the instructions of the manufacturers and as known to those skilled in the art. Total RNA was reverse transcribed (preferably using Superscript II RNaseH- Reverse Transcriptase, from Invitrogen, Karlsruhe, Germany) and subjected to Taqman analysis preferably using the Taqman 2xPCR Master Mix (from Applied Biosystems, Weiterstadt, Germany; the Mix contains according to the Manufacturer for example AmpliTaq Gold DNA Polymerase, AmpErase UNG, dNTPs with dUTP, passive reference Rox and optimized buffer components) on a GeneAmp 5700 Sequence Detection System (from Applied Biosystems, Weiterstadt, Germany).

Taqman analysis was performed preferably using the following primer/probe pairs:

For the amplification of Sac domain-containing inositol phosphatase 2 (sac2) (SEQ ID NO:2 ): 5'- CCT GGA TCG CAC CAA CG -3'; mouse sac2 reverse primer (SEQ ID NO:3 ): 5'- TTA AGC TGC TGT TCC ATG ACC A

-3'; Taqman probe (SEQ ID NO:4 ): (5/6-FAM) TCC AGG CTG CCA TAG CGC GC (5/6-TAMRA)

For the amplification of mouse solute carrier family 25 (mitochondrial carrier, Aralar) member 12 (Slc25a12) (SEQ ID NO:5 ): 5'- CCT GCC AAC CCT GAT CAC A -3'; mouse Slc25a12 reverse primer (SEQ ID NO:6 ): 5'- TTT CAA TGC CAG CGA AAG TG -3'; Taqman probe (SEQ ID NO:7 ): (5/6-FAM) CGG TGG CTA CAG ACT TGC CAC GG (5/6-TAMRA)

For the amplification of mouse solute carrier family 25 (mitochondrial carrier; adenine nucleotide translocator), member 13 (Slc25a13) (SEQ ID NO:8 ): 5'- AGC GGT GGT TCT ATG TCG ATT T -3'; mouse Slc25a13 reverse primer (SEQ ID NO:9 ): 5'- CGG GAT TTA GGA ACC GGC T -3'; Taqman probe (SEQ ID NO:10 ): (5/6-FAM) AGG CGT GAA GCC CGT GGG ATC T (5/6-TAMRA)

For the amplification of mouse myelin gene expression factor 2 (mef2) (SEQ ID NO:11 ): 5'- ACA AGG ATG GCA AGA GCA GAG -3'; mouse mef2 reverse primer (SEQ ID NO:12 ): 5'- ATG GAA ATT GCT TGG ACT GCT T -3'; Taqman probe (SEQ ID NO:13 ): (5/6-FAM) CAT GGG CAC TGT CAC TTT TGA GCA GG (5/6-TAMRA)

In the figures the relative RNA-expression is shown on the Y-axis. In Figures 4A and B, 8A, B, C, and D, and 16A, B, and C, the tissues tested are given on the X-axis. "WAT" refers to white adipose tissue, "BAT" refers to brown adipose tissue.

As shown in Figure 4A, real time PCR (Taqman) analysis of the expression of the Sac domain-containing inositol phosphatase 2 (SAC2) RNA in mammalian (mouse) tissues revealed that SAC2 is highly expressed in hypothalamus, brain, WAT, spleen and kidney. Figure 4B shows that SAC2 is upregulated in BAT and pancreas of fasted animals as well as ob / ob